MRC Holland Support

Support > Help Centre > MLPA & Coffalyser.Net > Experimental Setup > Sample Selection & Requirements > Is the purity and quality of the sample DNA important for (digital)MLPA?

Is the purity and quality of the sample DNA important for (digital)MLPA?

This article was retrieved from support.mrcholland.com on Saturday, 20th April 2024.

Sample purity is very important for conventional MLPA and digitalMLPA while sample quality is usually relatively unimportant for successful reactions.

Purity

Sample purity is of utmost importance for both conventional MLPA and digitalMLPA. Both techniques are based on a multiplex reaction, and have a higher sensitivity to impurities than some conventional PCR tests. Impurities such as ethanol, metal ions, salt, EDTA or phenol/Trizol remnants can decrease the activity of one of the enzymes or alter the properties of some probes, which can lead to poor or variable results.

For example, some probes are more sensitive to polymerase activity than others, which also makes them more sensitive to impurities that inhibit the polymerase. In conventional MLPA, longer amplicons are generally more sensitive to reduced polymerase activity than shorter fragments, which can lead to a characteristic sloping pattern.

The presence of NaCl or other ionic impurities, such as Mg²⁺, can also result in incomplete denaturation of the sample DNA. If the sample DNA is not completely denatured during the initial denaturation step in the protocol, probes may not be able to bind to their target sequences as efficiently. Probes that target GC-rich regions, which are more difficult to denature, will be more severely affected.

Read more about ways to improve the purity of DNA samples.

Quality

The quality of the DNA tends to have less of an effect on relative probe signals. For example, fragmentation does not strongly affect most probes because the target sequences are relatively short. Sample DNA that has degraded to the extent that a 1000 nt PCR is no longer possible may still produce good (digital)MLPA results.

Some chemical modifications of the DNA can have a more profound effect on (digital)MLPA results. Depurination, for example, affects probes differently depending on the amount and location of purines in the probes' target sequences. You can read

more about depurination and how to prevent it in this article.

Tags
digitalMLPA
MLPA

Related Pages

- Does RNA interfere with (digital)MLPA?
- What can I do about impurities in my DNA sample?
- <u>Is there a specific DNA extraction method I need to use for (digital)MLPA samples?</u>
- <u>Is it possible to use DNA extracted from formalin-fixed paraffin-embedded (FFPE)</u> <u>tissue for (digital)MLPA?</u>
- What are the sample and buffer requirements for (digital)MLPA?

Disclaimer

The information provided in this material is correct for the majority of our products. However, for certain applications, the instructions for use may differ. In the event of conflicting information, the relevant instructions for use take precedence.